

Diagnostic Parameters for Monitoring Peri-implant Conditions

Giovanni E. Salvi, Dr Med Dent¹/Niklaus P. Lang, Prof Dr Med Dent, Dr hc, FRCPS, PhD, MS²

Purpose: To review the literature on clinical, radiographic, and biochemical parameters used for monitoring peri-implant conditions. **Materials and Methods:** A MEDLINE search was conducted that included articles published in English until the end of August 2003. Results from human and experimental animal studies are presented. **Results:** The parameters that may be used to assess the presence of peri-implant health and the severity of peri-implant disease include plaque assessment, mucosal conditions, peri-implant probing depth, width of the peri-implant keratinized mucosa, peri-implant sulcus fluid analysis, suppuration, implant mobility and discomfort, resonance frequency analysis, and radiographic evaluation. **Discussion:** Based on the analysis of the available evidence, it appears reasonable to use a number of clinical and radiographic parameters to discriminate between peri-implant health and disease. **Conclusions:** Systematic and continuous monitoring of peri-implant tissues during maintenance care is recommended for the early diagnosis of peri-implant disease. *INT J ORAL MAXILLOFAC IMPLANTS* 2004;19(SUPPL):116–127

Key words: clinical parameters, dental implants, dental radiography, diagnosis, long-term evaluation, peri-implant disease

Oral endosseous implant systems with 2 different healing modalities (submerged and nonsubmerged) have been developed and used successfully for the rehabilitation of partially or completely edentulous patients.^{1–20} Knowledge of the biology of osseointegration and peri-implant soft tissue healing has expanded rapidly.^{21–27} A comparative study in the beagle dog²⁸ has provided histologic evidence that the peri-implant hard and soft tissues around 1-stage and 2-stage implant systems do not

significantly differ with respect to morphology and composition. Furthermore, studies have provided clinical and radiographic evidence that 2-part implant systems can successfully osseointegrate in the mandible when a nonsubmerged surgical protocol is applied for implant placement.^{29–32}

At the population level, longitudinal evaluation of oral implant systems is of primary importance for the assessment of long-term survival and complication rates of each system, for the determination of factors affecting the success of therapy, and for the identification of specific problems. At the individual level, clinical peri-implant evaluation is necessary for the detection of early signs of disease and for the planning of therapeutic interventions. An unbiased comparison of different implant systems is only meaningful if the stages of peri-implant disease are defined and if appropriate clinical parameters and indices are available.

As established in 1993 at the First European Workshop on Periodontology in Ittingen, Switzerland, *peri-implant disease* is a collective term for

¹Assistant Professor, Department of Periodontology and Fixed Prosthodontics, University of Berne, School of Dental Medicine, Berne, Switzerland.

²Chairman, Department of Periodontology and Fixed Prosthodontics, University of Berne, School of Dental Medicine, Berne, Switzerland.

Correspondence to: Dr Giovanni E. Salvi, University of Berne, School of Dental Medicine, Freiburgstrasse 7, CH-3010 Berne, Switzerland. Fax: +41-31-632-49-15. E-mail: giovanni.salvi@zmk.unibe.ch

Table 1 Indices Used to Assess Plaque Accumulation Around Oral Implants

Score	Mombelli et al ⁴⁰ (mPI)	Lindquist et al ⁴⁷
0	No detection of plaque	No visible plaque
1	Plaque only recognized by running a probe across the smooth marginal surface of the implant	Local plaque accumulation
2	Plaque can be seen by the naked eye	General plaque accumulation greater than 25%
3	Abundance of soft matter	

inflammatory processes in the tissues surrounding an implant.³³ *Peri-implant mucositis* was defined as a reversible inflammatory process in the soft tissues surrounding a functioning implant, whereas *peri-implantitis* is an inflammatory process additionally characterized by loss of peri-implant bone. A subgingival biofilm formation has been shown in animal experiments and clinical studies to be an important etiologic factor for the initiation of peri-implant inflammation and subsequent loss of marginal bone.^{34–37} In contrast to early implant losses, implant loss occurring during function may be the result of biologic processes characterized by clinical signs (eg, implant mobility) that emerge only when an advanced and possibly irreversible state of the disease has been reached. Therefore, the clinical and radiographic parameters routinely used to monitor oral implants during maintenance care should be of high sensitivity and/or specificity, should be easy to measure, and should yield reproducible data. The aim of this review article is to summarize current scientific evidence on the available diagnostic parameters for the longitudinal monitoring of oral implants.

MATERIALS AND METHODS

A search of Medline/PubMed was performed up to and including August 2003. The search was limited to human and experimental animal studies published in English. The following search terms were used: dental implants, peri-implant health, peri-implant disease, peri-implant mucositis, peri-implantitis, probing depth, bleeding on probing, dental plaque, peri-implant sulcus fluid, peri-implant keratinized mucosa, implant mobility, supuration, long-term evaluation, and dental radiography. The journals *Clinical Oral Implants Research*, *Journal of Clinical Periodontology*, *Journal of Periodontology*, and *The International Journal of Oral and Maxillofacial Implants* were searched by hand up to July 2003. The selection criteria included all levels

of available evidence: systematic reviews, randomized controlled clinical trials, controlled clinical trials, prospective and retrospective cohort studies, and case reports of human and experimental animal studies. One reviewer (GES) screened titles and abstracts of the search results. The full text of publications of potential relevance was then obtained.

EVALUATION OF THE ORAL HYGIENE STANDARD

Plaque Assessment

Microbial biofilms have been shown to form on inert biomaterial surfaces in an aqueous environment.³⁸ Implants placed in the oral cavity represent artificial surfaces colonized by bacteria from saliva and ecologic niches such as periodontal pockets, tonsils, and crypts of the tongue. Experimental and human studies have provided evidence that formation and development of a microbial biofilm represents an important etiologic factor in the pathogenesis of peri-implant disease.^{34,37,39–42} Several microbiologic features of the subgingival biofilm around implants have been correlated with the presence of clinically detectable plaque.⁴³ Furthermore, periodontal pathogens from residual pockets of remaining teeth in patients treated for periodontal disease have been documented to colonize oral implants.^{44,45} Mombelli and coworkers⁴⁰ modified the original Plaque Index introduced by Silness and L e⁴⁶ to assess biofilm formation in the marginal area around ITI implants (mPI) (Institut Straumann, Waldenburg, Switzerland). Lindquist and associates⁴⁷ assessed oral hygiene levels according to a 3-point scale and reported a significant relationship between oral hygiene and peri-implant bone resorption over an observation period of 6 years. Therefore, it appears meaningful to monitor oral hygiene habits by quantifying plaque accumulation. Indices used to assess plaque accumulation around oral implants are presented in Table 1.

Table 2 Indices Used to Assess Marginal Mucosal Conditions Around Oral Implants

Score	Mombelli et al ⁴⁰ (mGI)	Apse et al ⁵⁰
0	No bleeding when a periodontal probe is passed along the mucosal margin adjacent to the implant	Normal mucosa
1	Isolated bleeding spots visible	Minimal inflammation with color change and minor edema
2	Blood forms a confluent red line on mucosal margin	Moderate inflammation with redness, edema, and glazing
3	Heavy or profuse bleeding	Severe inflammation with redness, edema, ulceration, and spontaneous bleeding without probing

EVALUATION OF THE PERI-IMPLANT MARGINAL TISSUES

Mucosal Conditions

In addition to redness and swelling of the marginal tissues, bleeding on probing (BOP), pocket formation, and suppuration have been reported to result from peri-implant infections.^{40,48} Assessment of these clinical signs has been considered important in the diagnosis of periodontal diseases. Therefore, the definition of peri-implant parameters based on periodontal indices such as the Gingival Index System⁴⁹ (GI) seems indicated. The GI⁴⁹ has been modified and adapted (mGI) for application around oral implants,⁴⁰ while a simplified GI has been proposed by Apse and associates.⁵⁰ Indices used to assess marginal mucosal conditions around oral implants are presented in Table 2.

Around implants, however, soft tissue texture and color depend on the normal appearance of the recipient tissues before implant placement, and may be influenced by the material characteristics of the implant surface.⁵¹ Furthermore, difficulties in recording mucosal inflammation have been reported, such as nonkeratinized peri-implant mucosa normally appearing redder than keratinized tissue.⁵² In a longitudinal study, only a weak correlation between GI scores and changes in peri-implant crestal bone level was reported.⁵²

Presence or Absence of Bleeding

BOP (notated in clinical records as BOP+) elicited after the insertion of a probe into the sulcus with light pressure (ie, 0.25 N) has been shown to detect the presence of an inflammatory lesion in the gingiva around teeth with a normal⁵³ and a healthy but reduced periodontium.⁵⁴ On the other hand, absence of bleeding on probing (BOP-) has been reported to represent periodontal health with a negative predictive value of 98.5%.^{55,56} BOP has been

used to assess peri-implant tissue conditions around implants. Lekholm and colleagues⁵⁷ found no correlation between BOP and histologic, microbiologic, or radiographic changes around implants. These authors hypothesized that bleeding could have been caused by inappropriate force transmission from the periodontal probe tip to the peri-implant soft tissues. These preliminary findings were confirmed in an animal study.⁵⁸ Conversely, findings from animal experiments using ITI implants yielded completely different results.⁵⁹ Healthy sites were characterized by absence of bleeding (0%), whereas both peri-implant mucositis and peri-implantitis sites showed substantially increased BOP (67% and 91%, respectively). The reason for these results might be attributed to the different probing forces applied by the various investigators. These findings were confirmed in a prospective study where absence of BOP had a high negative predictive value, thus serving as a predictor for stable peri-implant conditions.⁶⁰ Luterbacher and coworkers⁶¹ evaluated the diagnostic characteristics of different BOP prevalences alone or in combination with a microbiologic test for monitoring periodontal and peri-implant soft tissue conditions during maintenance therapy. The authors reported that BOP alone yielded higher diagnostic accuracy at implant sites compared with tooth sites. Furthermore, when combining positive microbiologic test and BOP of 75% or more, the positive predictive values were greater for implants than for teeth. With predictive values of 100% at BOP frequencies of 50% or more at implant sites, this parameter appears to play a central role in monitoring changes in peri-implant tissue conditions.

Peri-implant Probing Depth

In contrast to natural teeth, for which average periodontal probing depth (PD) has been reported, the physiologic depth of the peri-implant sulcus of successfully osseointegrated implants has been a matter

of debate. Increasing periodontal PD and loss of clinical attachment are pathognomonic for periodontal diseases. Pocket probing is therefore an important diagnostic process for the assessment of periodontal status and for the evaluation of periodontal therapy. The extent of probe penetration is influenced by factors such as probing force and angulation, probe tip diameter, roughness of the implant or root surface, inflammatory state of the periodontium, and firmness of the marginal tissues. Furthermore, it has to be realized that PD measurements may be affected by compromised access. Data demonstrate that the periodontal probe often fails to locate the histologic level of the connective tissue attachment around teeth.^{62,63}

The differences with respect to soft tissue composition, organization, and attachment between the gingiva and the root surface on the one hand and between the peri-implant mucosa and the implant surface on the other make the conditions for PD measurements around teeth and implants not fully comparable.^{37,64,65} One potential explanation influencing the differences in probe penetration is that most collagen fibers in the supracrestal connective tissue compartment have been demonstrated to run mostly in a parallel direction to the implant axis.^{51,66}

Ericsson and Lindhe⁵⁸ used a beagle dog model to compare the extent of probe penetration around teeth and implants under healthy soft tissue conditions. Compared with natural teeth, probe tip penetration around 2-stage submerged implants ended closer to the alveolar crest. The extent of peri-implant probe penetration has also been investigated around 1-stage oral implants in beagle dogs.⁵⁹ It was reported that density of the peri-implant tissues influences probe penetration. In the presence of inflamed peri-implant tissues, periodontal probes penetrate close to the alveolar bone, exceeding the connective tissue level by a mean of 0.52 mm. However, if healthy peri-implant conditions or mucositis are present, the probe tips may identify the histologic level of the supracrestal connective tissue attachment. One potential explanation for the different outcomes between the 2 above-mentioned studies may be attributed to the different probing forces employed (0.5 N versus 0.2 N). A recent experimental study in monkeys⁶⁵ has documented that PD measurements around teeth and implants are different. While no difference was observed with respect to probe penetration under healthy peri-implant/periodontal conditions, mild and severe marginal inflammation around implants was associated with a significantly deeper probe penetration into the supracrestal connective tissue when compared to that around teeth.

The magnitude of probe penetration into a periodontal pocket depends on the force applica-

tion.^{67,68} Furthermore, simultaneous recordings of probing PD and probing force before and after periodontal therapy have revealed that the force range chosen for repeated probing influences the amount of attachment level change determined.^{69,70} The tissue resistance to probing and the accuracy of depth measurement at different force levels (eg, 0.25, 0.5, 0.75, 1.0, and 1.25 N) were compared around nonsubmerged ITI dental implants and teeth in 11 subjects.⁷¹ It was concluded that peri-implant PD measurements are more sensitive to force variation than the corresponding measurements around teeth.

Correlations between bone levels recorded on radiographs and the extent of peri-implant probe penetration have been observed. In the case of screw-type implants, the probe tip appeared to stop 1.4 mm coronally to the bone level.⁷² The mean discrepancy between probe penetration and the location of the bone margin in radiographs was 1.17 mm in 100 nonsubmerged hollow-screw and hollow-cylinder implants measured 1 year after implantation.⁷³ In general, studies have indicated that successful implants allow probe penetration of approximately 3 mm.^{40,50,73-78}

Implant shape and surface texture influence penetration of the probe tip. Peri-implant probing is impossible around some implant systems because of characteristics of the design (eg, concavities, shoulders, suprastructure, or steps). Lack of surface smoothness, as with plasma-coated, sandblasted, acid-etched, or threaded implants, might interfere with probe penetration when bone resorption has reached this level and may lead to underestimation of pocket depth.⁷⁹ Although convincing evidence is lacking, some authors have expressed concerns about the possibility of introducing bacteria into the peri-implant tissues and about damaging the implant surface with a metallic periodontal probe.^{58,65,77,79} Other authors concluded that increased pocket depth could be correlated with a higher degree of inflammation of the peri-implant mucosa^{41,57,72,80} but not necessarily with peri-implant bone loss.^{81,82} However, absolute values of PD have to be interpreted in the context of surgical implant positioning, eg, submucosal implant placement in esthetic anterior sites versus conventional implant placement in posterior non-esthetic sites. Progressive increases in PD may be an alarming sign. Therefore, the establishment of baseline PD values at the time of delivery of the prosthetic suprastructure is of critical importance in allowing comparison with future PD measurements.

Peri-implant probing should also include the location of the soft tissue margin relative to a fixed landmark point on the implant (eg, implant shoulder for

1-stage nonsubmerged implant systems) or its supra-structure. The combined full-mouth PI and the BOP scores, as well as the mean attachment level of all teeth in a group of partially edentulous subjects, were shown to significantly influence mean PD and connective tissue level around implants.⁸³ If peri-implantitis is associated with marginal recession, then PD alone may not accurately reflect peri-implant bone loss, whereas increasing loss of connective tissue level is a definite sign of peri-implant pathology. In a longitudinal study, Bragger and coworkers⁸⁴ found that the connective tissue level in combination with the radiographic parameters obtained 2 years after implant loading were good predictors of the peri-implant tissue condition. Repeated peri-implant PD measurements may be performed with higher reproducibility by means of an automated controlled-force periodontal probe.⁸⁵ Furthermore, animal experiments have shown that, as is the case around teeth,⁸⁶ peri-implant probing disrupts the epithelial attachment but does not cause permanent damage to the transmucosal soft tissue seal. In one study, complete attachment of the junctional epithelium was re-established after 5 days following probing using a conventional periodontal probe.⁸⁷

Width of Peri-implant Keratinized Mucosa

Clinical and experimental studies^{88,89} have failed to support the concept of an “adequate width” of keratinized tissue adjacent to natural teeth for the maintenance of periodontal health. Implant research has also focused on the necessity of the presence of keratinized mucosa around oral implants. No differences in peri-implant soft tissue recession or bone loss have been found between sites with or without keratinized mucosa following plaque-induced breakdown at implants placed in dogs.⁹⁰ On the other hand, ligated titanium or hydroxyapatite-coated implants in monkeys with minimal or no keratinized mucosa demonstrated significantly more recession and connective tissue loss than those surrounded by keratinized tissue.^{91,92} This suggests that the absence of keratinized mucosa around implants seems to increase the susceptibility of plaque-induced peri-implant tissue destruction. These findings have been confirmed in other studies,^{83,93} suggesting that the presence of keratinized mucosa around implants is strongly correlated with optimal soft and hard tissue health. However, longitudinal clinical studies have failed to reveal major differences in the progression of lesions around implants placed in sites with or without keratinized mucosa, or that the lack of an attached portion of masticatory mucosa may jeopardize the maintenance of soft tissue health.^{74,94–97} Furthermore, in

the presence of good oral hygiene, the nature of the mucosa may have little influence on the long-term survival of implants. However, suboptimal oral hygiene may lead to greater tissue damage around implants within alveolar mucosa than around implants within keratinized tissue. Proper oral hygiene procedures may also be facilitated in the presence of an adequate band of keratinized mucosa. Prospective longitudinal controlled clinical trials will have to be performed to further elucidate the potential role of a sealing effect of keratinized mucosa on long-term peri-implant health.

Peri-implant Sulcus Fluid Analysis

Several biochemical mediators in the gingival crevicular fluid (GCF) around natural teeth have been identified as potential host markers for periodontal disease activity and progression.⁹⁸ To date, only a few studies have reported on the association between signs of peri-implant inflammation and increased levels of inflammatory mediators in the peri-implant sulcus fluid (PISF). A pilot study⁹⁹ investigated whether the crevicular fluid volume around osseointegrated implants shows a relationship to peri-implant soft tissue condition. The results indicated a close relationship between PISF volume and plaque accumulation as well as degree of peri-implant soft tissue inflammation. Numerous investigations of potential diagnostic markers of stable and diseased peri-implant conditions have focused on the sulcus fluid analysis of several mediators, including protease activity^{60,100,101}; collagenase, gelatinase, and elastase activity^{102–107}; aspartate aminotransferase¹⁰⁸; glycosaminoglycans^{109,110}; and proinflammatory mediators such as interleukin-1 β (IL-1 β) and prostaglandin E₂ (PGE₂).^{80,111} Kao and coworkers¹¹¹ reported that PISF-IL-1 β levels around diseased implants were approximately 3 times higher than those around stable implants, thus providing evidence for the involvement of this catabolic cytokine in peri-implant bone destruction.

In a 3-year longitudinal investigation, Behneke and associates⁸² were able to show a positive correlation between PISF volume and the amount of bone resorption. In a subsequent report of the 5-year data,² the percentage of sites exhibiting elevated PISF rates increased significantly in the second half of the observation period. Using a cross-sectional study design, Salcetti and coworkers⁸⁰ investigated the production of IL-1 β , PGE₂, interleukin-6, platelet-derived growth factor, and transforming growth factor beta in the PISF of patients with 1 or more failing titanium implants. Several of these patients had at least 1 other stable implant that did not present with clinical signs of

inflammation or radiographic evidence of peri-implant bone loss. Significant elevations in PISF levels of IL-1 β , PGE₂, and platelet-derived growth factor in subjects with failing implant sites were detected when compared with patients with healthy control implants. The significant elevations of IL-1 β and PGE₂ at both failing implant sites and at stable implant sites in the same subject indicate that an increased local host response is detectable at the patient level as well as at local sites of peri-implant inflammation.

The finding that enzymes from polymorphonuclear granulocytes (PMN) are detected in high concentrations at sites with peri-implantitis may indicate enhanced PMN cell activity.^{105,107} Hultin and colleagues¹⁰⁵ analyzed the composition of PISF at implants in patients with “stable marginal tissue conditions” and peri-implantitis. Implant sites with peri-implantitis had higher concentrations of lactoferrin and elastase activity than control sites. Similarly, Plagnat and associates¹⁰⁷ collected and analyzed PISF from sites with and without clinical and radiographic signs of peri-implantitis. The authors reported that PISF levels of elastase, alpha2-macroglobulin, and alkaline phosphatase were significantly higher at diseased sites than at healthy sites, and that the levels of these markers correlated with clinical symptoms. On the other hand, similar low levels of the above-mentioned markers were found both at baseline and at the 3-year examination in the sulcus fluid around successful implants placed in esthetic anterior sites, indicating stable biochemical peri-implant conditions.¹⁰⁴ Collectively, these data document an important implication of catabolic inflammatory mediators in peri-implant tissue breakdown and indicate a potential value of biochemical markers for monitoring the host response during the supportive phase of implant therapy.

Suppuration

High numbers of PMN cells have been detected around implants that are associated with severe signs of mucosal inflammation.⁴⁸ Several histopathologic and immunohistochemical analyses of tissues surrounding implants with signs of peri-implantitis, ie, clinical signs of inflammation and advanced bone loss, have revealed the presence of large inflammatory cell infiltrates.^{35,112–115} Sanz and associates¹¹⁵ analyzed soft tissue biopsies from patients with peri-implantitis and reported that a considerable portion of the connective tissue (ie, 65%) was occupied by an inflammatory infiltrate. Esposito and coworkers¹¹² analyzed the characteristics of soft tissues surrounding failing implants

immunohistochemically. They reported that the marginal portion of the specimens was characterized by an “intense inflammatory and immunologic response.” Piattelli and colleagues¹¹⁴ described histopathologic characteristics of 230 retrieved implants. The authors reported that around implants removed because of peri-implantitis, “an inflammatory infiltrate composed of macrophages, lymphocytes, and plasma cells was observed in the connective tissue.” Gualini and Berglundh¹¹³ reported that peri-implantitis lesions contained significantly greater proportions of B-lymphocytes and PMN cells than mucositis lesions. Collectively, the observation that large numbers of inflammatory cells, including PMN cells, occupy the connective tissue infiltrate may explain the presence of suppuration at sites with advanced peri-implant disease.

EVALUATION OF THE BONE-IMPLANT INTERFACE

Implant Mobility and Discomfort

Primary stability at the time of implant placement has been recognized as an important prerequisite for the achievement of osseointegration.^{74,94,116} The establishment and maintenance of direct contact at the bone-implant interface are requirements for long-term implant success. Implant mobility is an indication of lack of osseointegration. Even if peri-implant disease has progressed relatively far, implants may still appear immobile because of some residual direct bone-to-implant contact. The recording of implant mobility may be a very specific—but not at all sensitive—clinical parameter in detecting loss of osseointegration. This parameter more likely detects the final stage of osseo-disintegration and, therefore, represents a late implant loss. Furthermore, pain or discomfort may be associated with increased implant mobility and could be one of the first signs indicating a failing implant.^{117,118} Persistent discomfort may be evident long before any radiographic change is detectable.¹¹⁹

Longitudinal assessment of individual implant mobility may be performed for screw-retained suprastructures. For obvious reasons, this cannot be applied to all cemented and tooth/implant-supported restorations. An electronic device (Periotest; Siemens, Bensheim, Germany), originally designed to measure the damping characteristics of the periodontium around natural teeth,^{120,121} has been recommended to monitor initial degrees of implant mobility or horizontal displacement. However, differences in Periotest values (PTVs) have been reported for implants in the mandible and in the

maxilla,⁷³ with implants in the maxilla showing significantly higher PTVs. In patients treated with Brånemark System implants (Nobel Biocare, Göteborg, Sweden), this procedure was found to be related to the type of jaw treated, implant and abutment length, condition of the peri-implant tissues, and bone density.¹²² Despite some positive claims for this method,^{2,82,123,124} the prognostic accuracy of PTVs for the diagnosis of peri-implantitis and early signs of implant failure has been criticized because of the lack of resolution, poor sensitivity, and susceptibility to operator variables.¹²⁵

Resonance Frequency Analysis

A new, noninvasive device based on the principles of resonance frequency analysis (RFA) has been developed to measure primary implant stability and to monitor implant stability over time.¹²⁶ This method evaluates the stiffness of the bone-implant interface by means of a signal transducer connected to a frequency response analyzer (Osstell; Integration Diagnostics, Göteborg, Sweden). The resonance frequency of the transducer-implant unit is calculated from the peak amplitude of the signal and is graphically illustrated on the Osstell display as the peak of a frequency-amplitude plot. In addition, an implant stability quotient (ISQ) is displayed as a number between 1 and 100. This ISQ value has been introduced to quantify the frequency measurements of oral implants with a range between 3,500 and 8,500 Hz. Several investigations¹²⁷⁻¹³² have shown that the ISQ value of a stable osseointegrated implant increases with time, suggesting an increase in the bone-implant contact area. On the other hand, crestal bone loss around implants has been correlated with loss of implant stability.^{124,129} This may allow detection of an increase in implant mobility before clinical signs are recorded.¹³³ However, conclusive data on the bone-implant interface and RFA values are still lacking.¹³⁴

RADIOGRAPHIC EVALUATION

Long-term preservation of crestal bone height around osseointegrated implants is often used as a primary success criterion for different implant systems. Originally, a mean crestal bone loss ≥ 1.5 mm during the first year after loading and ≥ 0.2 mm/year thereafter had been proposed as one of the major success criteria.^{94,135} This success criterion, however, has recently been questioned, because longitudinal studies have provided evidence that crestal bone loss around osseointegrated implants in well-maintained patients may be minimal.^{2,18,136}

Conventional radiography represents a widely accepted technique for the long-term evaluation of marginal bone changes at interproximal sites of osseointegrated implants. In general, the long-cone paralleling technique, supported by positioning devices, is used. It should be noted that conventional radiography yields a high proportion of false negative findings, ie, it has low sensitivity in the detection of early pathologic and/or bone remodeling changes.¹³⁷ Therefore, radiographic methods are confirmatory rather than exploratory and should only be considered in conjunction with assessment of the clinical parameters.¹³⁸ Nevertheless, the distance from a landmark on the implant (eg, implant shoulder for 1-stage transmucosal implant systems or apical termination of the cylindrical portion of the implant for 2-stage submerged implant systems) to the alveolar bone crest represents a reliable parameter for long-term monitoring in clinical practice.

It should be pointed out that radiographic evidence of bone-to-implant contact does not imply osseointegration on a histologic level.¹³⁹ More importantly, if clinical parameters indicate peri-implant disease (eg, increased PD, BOP+, suppuration), additional radiographs should be obtained to evaluate the extent of peri-implant crestal bone loss. For longitudinal clinical research purposes, radiographs should be obtained at baseline and at 1-, 3-, and 5-year intervals. Thereafter, they should be obtained every 5 years if marginal peri-implant bone stability has been demonstrated.¹⁴⁰

Computer-assisted image analysis has been shown to improve the diagnostic accuracy (ie, increased sensitivity) of detecting minimal periodontal tissue changes.^{137,141} Consequently, the use of digital image analysis has expanded into implant dentistry to monitor peri-implant bone healing and gain or loss of alveolar bone density.¹⁴²

CONCLUSION

Evidence from the presented literature indicates that the use of a number of clinical, biochemical, and radiographic parameters is meaningful in the evaluation of peri-implant tissue status. Research efforts are currently under way to relate biologic parameters to morphologic changes in peri-implant structures. However, reliable prognostic indicators for peri-implant hard and soft tissue changes are still lacking. This is not surprising, because the same phenomenon (ie, development of diagnostic tests for the assessment of active periodontal tissue destruction) has challenged periodontal research in recent decades.

ACKNOWLEDGMENT

This study was supported by the Clinical Research Foundation (CRF) for the Promotion of Oral Health, University of Berne, Switzerland.

REFERENCES

- Behneke A, Behneke N, d'Hoedt B. The longitudinal clinical effectiveness of ITI solid-screw implants in partially edentulous patients: A 5-year follow-up report. *Int J Oral Maxillofac Implants* 2000;15:633–645.
- Behneke A, Behneke N, d'Hoedt B. A 5-year longitudinal study of the clinical effectiveness of ITI solid-screw implants in the treatment of mandibular edentulism. *Int J Oral Maxillofac Implants* 2002;17:799–810.
- Brägger U, Aeschlimann S, Bürgin W, Hämmerle CHF, Lang NP. Biological and technical complications and failures with fixed partial dentures (FPD) on implants and teeth after four to five years of function. *Clin Oral Implants Res* 2001;12:26–34.
- Lambrecht JT, Filippi A, Kunzel AR, Schiel HJ. Long-term evaluation of submerged and nonsubmerged ITI solid-screw titanium implants: A 10-year life table analysis of 468 implants. *Int J Oral Maxillofac Implants* 2003;18:826–834.
- Brocard D, Barthet P, Baysse E, et al. A multicenter report on 1,022 consecutively placed ITI implants: A 7-year longitudinal study. *Int J Oral Maxillofac Implants* 2000;15:691–700.
- Buser D, Mericske-Stern R, Dula K, Lang NP. Clinical experience with one-stage, non-submerged dental implants. *Adv Dent Res* 1999;13:153–161.
- Ferrigno N, Laureti M, Fanali S, Grippaudo G. A long-term follow-up study of non-submerged ITI implants in the treatment of totally edentulous jaws. Part I: Ten-year life table analysis of a prospective multicenter study with 1286 implants. *Clin Oral Implants Res* 2002;13:260–273.
- Hardt CRE, Gröndahl K, Lekholm U, Wennström JL. Outcome of implant therapy in relation to experienced loss of periodontal bone support. A retrospective 5-year study. *Clin Oral Implants Res* 2002;13:488–494.
- Karoussis IK, Brägger U, Salvi GE, Bürgin W, Lang NP. Effect of implant design on survival and success rates of titanium oral implants: A 10-year prospective cohort study of the ITI Dental Implant System. *Clin Oral Implants Res* 2004;15:8–17.
- Karoussis IK, Müller S, Salvi GE, Heitz-Mayfield LJA, Brägger U, Lang NP. Association between periodontal and periimplant conditions: A 10-year prospective cohort study of the ITI Dental Implant System. *Clin Oral Implants Res* 2004;15:1–7.
- Karoussis IK, Salvi GE, Heitz-Mayfield LJA, Brägger U, Hämmerle CHF, Lang NP. Long-term implant prognosis in patients with and without a history of chronic periodontitis: A 10-year prospective cohort study of the ITI Dental Implant System. *Clin Oral Implants Res* 2003;14:329–339.
- Levine RA, Clem D, Beagle J, et al. Multicenter retrospective analysis of the solid-screw ITI implant for posterior single-tooth replacements. *Int J Oral Maxillofac Implants* 2002;17:550–556.
- Mericske-Stern R, Aerni D, Geering AH, Buser D. Long-term evaluation of non-submerged hollow-cylinder implants. Clinical and radiographic results. *Clin Oral Implants Res* 2001;12:252–259.
- Mericske-Stern R, Grütter L, Rösch R, Mericske E. Clinical evaluation and prosthetic complications of single tooth replacements by non-submerged implants. *Clin Oral Implants Res* 2001;12:309–318.
- Moberg LE, Kondell PA, Kullmann L, Heimdahl A, Gynther GW. Evaluation of single-tooth restorations on ITI dental implants. A prospective study of 29 patients. *Clin Oral Implants Res* 1999;10:45–53.
- Moberg LE, Kondell PA, Sagulin GB, Bolin A, Heimdahl A, Gynther GW. Brånemark System and ITI Dental Implant System for treatment of mandibular edentulism. A comparative randomized study: 3-year follow-up. *Clin Oral Implants Res* 2001;12:450–461.
- Naert IE, Koutsikakis G, Duyck J, Quirynen M, Jacobs R, van Steenberghe D. Biologic outcome of implant-supported restorations in the treatment of partial edentulism. Part I: A longitudinal clinical evaluation. *Clin Oral Implants Res* 2002;13:381–389.
- Naert IE, Koutsikakis G, Quirynen M, Duyck J, van Steenberghe D, Jacobs R. Biologic outcome of implant-supported restorations in the treatment of partial edentulism. Part II: A longitudinal radiographic study. *Clin Oral Implants Res* 2002;13:390–395.
- Romeo E, Chiapasco M, Ghisolfi M, Vogel G. Long-term clinical effectiveness of oral implants in the treatment of partial edentulism. Seven-year life table analysis of a prospective study with the ITI Dental Implant System used for single-tooth restorations. *Clin Oral Implants Res* 2002;13:133–143.
- Weber H-P, Crohin CC, Fiorellini JP. A 5-year prospective clinical and radiographic study of non-submerged dental implants. *Clin Oral Implants Res* 2000;11:144–153.
- Abrahamsson I, Zitzmann NU, Berglundh T, Linder E, Wennerberg A, Lindhe J. The mucosal attachment to titanium implants with different surface characteristics. An experimental study in dogs. *J Clin Periodontol* 2002;29:448–455.
- Berglundh T, Abrahamsson I, Lang NP, Lindhe J. De novo alveolar bone formation adjacent to endosseous implants. A model study in the dog. *Clin Oral Implants Res* 2003;14:251–262.
- Botticelli D, Berglundh T, Buser D, Lindhe J. Appositional bone formation in marginal defects at implants. An experimental study in the dog. *Clin Oral Implants Res* 2003;14:1–9.
- Botticelli D, Berglundh T, Buser D, Lindhe J. The jumping distance revisited. An experimental study in the dog. *Clin Oral Implants Res* 2003;14:35–42.
- Moon I-S, Berglundh T, Abrahamsson I, Linder E, Lindhe J. The barrier between the keratinized mucosa and the dental implant. An experimental study in the dog. *J Clin Periodontol* 1999;26:658–663.
- Wennerberg A, Sennerby L, Kultje C, Lekholm U. Some soft tissue characteristics at implant abutments with different topography. A study in humans. *J Clin Periodontol* 2003;30:88–94.
- Zitzmann NU, Berglundh T, Marinello CP, Lindhe J. Expression of endothelial adhesion molecules in the alveolar ridge mucosa, gingiva and periimplant mucosa. *J Clin Periodontol* 2002;29:490–495.
- Abrahamsson I, Berglundh T, Wennström J, Lindhe J. The periimplant hard and soft tissues at different implant systems. A comparative study in the dog. *Clin Oral Implants Res* 1996;7:212–219.

29. Bernard JP, Belser UC, Martinet JP, Borgis SA. Osseointegration of Brånemark fixtures using a single-step operating technique. A preliminary prospective one-year study in the edentulous mandible. *Clin Oral Implants Res* 1995;6:122–129.
30. Ericsson I, Randow K, Glantz P-O, Lindhe J, Nilner K. Clinical and radiographical features of submerged and non-submerged titanium implants. *Clin Oral Implants Res* 1994;5:185–189.
31. Heydenrijk K, Raghoobar GM, Meijer HJA, van der Reijden WA, van Winkelhoff A-J, Stegenga B. Two-part implants inserted in a one-stage or two-stage procedure. A prospective comparative study. *J Clin Periodontol* 2002;29:900–909.
32. Heydenrijk K, Raghoobar GM, Meijer HJA, van der Reijden WA, van Winkelhoff A-J, Stegenga B. Two-stage IMZ implants and ITI implants inserted in a single-stage procedure. A prospective comparative study. *Clin Oral Implants Res* 2002;13:371–380.
33. Albrektsson T, Isidor F. Consensus report of session IV. In: Lang NP, Karring T (eds). *Proceedings of the 1st European Workshop on Periodontology*. London: Quintessence, 1994:365–369.
34. Lang NP, Brägger U, Walther D, Beamer B, Kornman KS. Ligature-induced periimplant infection in cynomolgus monkeys. I. Clinical and radiographic findings. *Clin Oral Implants Res* 1993;4:2–11.
35. Lindhe J, Berglundh T, Ericsson I, Liljenberg B, Marinello C. Experimental breakdown of periimplant and periodontal tissues. A study in the beagle dog. *Clin Oral Implants Res* 1992;3:9–16.
36. Schou S, Holmstrup P, Reibel J, Juhl M, Hjørting-Hansen E, Kornman KS. Ligature-induced marginal inflammation around osseointegrated implants and ankylosed teeth: Stereologic and histologic observations in cynomolgus monkeys (*Macaca fascicularis*). *J Periodontol* 1993;64:529–537.
37. Schou S, Holmstrup P, Stoltze K, Hjørting-Hansen E, Kornman KS. Ligature-induced marginal inflammation around osseointegrated implants and ankylosed teeth. Clinical and radiographic observations in cynomolgus monkeys (*Macaca fascicularis*). *Clin Oral Implants Res* 1993;4:12–22.
38. Gristina AG. Biomaterial-centered infection: Microbial adhesion versus tissue integration. *Science* 1987;237:1588–1595.
39. Berglundh T, Lindhe J, Ericsson I, Marinello CP, Liljenberg B. Soft tissue reactions to de novo plaque formation on implants and teeth. An experimental study in the dog. *Clin Oral Implants Res* 1992;3:1–8.
40. Mombelli A, Van Oosten MAC, Schürch E, Lang NP. The microbiota associated with successful or failing osseointegrated titanium implants. *Oral Microbiol Immunol* 1987;2:145–151.
41. Pontoriero R, Tonelli MP, Carnevale G, Mombelli A, Nyman SR, Lang NP. Experimentally induced periimplant mucositis. A clinical study in humans. *Clin Oral Implants Res* 1994;5:254–259.
42. Zitzmann NU, Abrahamsson I, Berglundh T, Lindhe J. Soft tissue reactions to plaque formation at implant abutments with different surface topography. An experimental study in dogs. *J Clin Periodontol* 2002;29:456–461.
43. Mombelli A, Mericske-Stern R. Microbiological features of stable osseointegrated implants used as abutments for overdentures. *Clin Oral Implants Res* 1990;1:1–7.
44. Lee KH, Maiden MFJ, Tanner ACR, Weber HP. Microbiota of successful osseointegrated dental implants. *J Periodontol* 1999;70:131–138.
45. Mombelli A, Marxer M, Gaberthüel T, Grunder U, Lang NP. The microbiota of osseointegrated implants in patients with a history of periodontal disease. *J Clin Periodontol* 1995;22:124–130.
46. Silness J, Løe H. Periodontal disease in pregnancy. II. Correlation between oral hygiene and periodontal condition. *Acta Odontol Scand* 1964;22:121–135.
47. Lindquist LW, Rocker B, Carlsson GE. Bone resorption around fixtures in edentulous patients treated with mandibular fixed tissue-integrated prostheses. *J Prosthet Dent* 1988;59:59–63.
48. Rams TE, Roberts TW, Tatum H Jr, Keyes PH. The subgingival microbial flora associated with human dental implants. *J Prosthet Dent* 1984;51:529–534.
49. Løe H. The Gingival Index, the Plaque Index and the Retention Index Systems. *J Periodontol* 1967;38(suppl):610–616.
50. Apse P, Zarb GA, Schmitt A, Lewis DW. The longitudinal effectiveness of osseointegrated dental implants. The Toronto study: Periimplant mucosal response. *Int J Periodontics Restorative Dent* 1991;11:95–111.
51. Listgarten MA, Lang NP, Schroeder HE, Schroeder A. Periodontal tissues and their counterparts around endosseous implants. *Clin Oral Implants Res* 1991;2:1–19.
52. Chaytor DV. The longitudinal effectiveness of osseointegrated dental implants. The Toronto study: Bone level changes. *Int J Periodontics Restorative Dent* 1991;11:113–125.
53. Lang NP, Nyman S, Senn C, Joss A. Bleeding on probing as it relates to probing pressure and gingival health. *J Clin Periodontol* 1991;18:257–261.
54. Karayiannis A, Lang NP, Joss A, Nyman S. Bleeding on probing as it relates to probing pressures and gingival health in patients with a reduced but healthy periodontium. A clinical study. *J Clin Periodontol* 1991;19:471–475.
55. Joss A, Adler R, Lang NP. Bleeding on probing. A parameter for monitoring periodontal conditions in clinical practice. *J Clin Periodontol* 1994;21:402–408.
56. Lang NP, Adler R, Joss A, Nyman S. Absence of bleeding on probing. An indicator of periodontal stability. *J Clin Periodontol* 1990;17:714–721.
57. Lekholm U, Eriksson A, Adell R, Slots J. The conditions of the soft tissues at tooth and fixture abutments supporting fixed bridges. A microbiological and histological study. *J Clin Periodontol* 1986;13:558–562.
58. Ericsson I, Lindhe J. Probing depth at implants and teeth. An experimental study in the dog. *J Clin Periodontol* 1993;20:623–627.
59. Lang NP, Wetzel AC, Stich H, Caffesse RG. Histologic probe penetration in healthy and inflamed periimplant tissues. *Clin Oral Implants Res* 1994;5:191–201.
60. Jepsen S, Rühling A, Jepsen K, Ohlenbusch B, Albers H-K. Progressive periimplantitis. Incidence and prediction of periimplant attachment loss. *Clin Oral Implants Res* 1996;7:133–142.
61. Luterbacher S, Mayfield L, Brägger U, Lang NP. Diagnostic characteristics of clinical and microbiological tests for monitoring periodontal and periimplant mucosal tissue conditions during supportive periodontal therapy (SPT). *Clin Oral Implants Res* 2000;11:521–529.
62. Armitage GC, Svanberg GK, Løe H. Microscopic evaluation of clinical measurements of connective tissue attachment levels. *J Clin Periodontol* 1977;4:173–190.
63. Magnusson I, Listgarten MA. Histological evaluation of probing depth following periodontal treatment. *J Clin Periodontol* 1980;7:26–31.

64. Berglundh T, Lindhe J, Ericsson I, Marinello CP, Liljenberg B, Thomsen P. The soft tissue barrier at implants and teeth. *Clin Oral Implants Res* 1991;2:81–90.
65. Schou S, Holmstrup P, Stoltze K, Hjørting-Hansen E, Fiehn NE, Skovgaard LT. Probing around implants and teeth with healthy or inflamed peri-implant mucosa/gingiva. A histologic comparison in cynomolgus monkeys (*Macaca fascicularis*). *Clin Oral Implants Res* 2002;13:113–126.
66. Listgarten MA, Lai CH. Ultrastructure of the intact interface between an endosseous epoxy resin dental implant and the host tissues. *J Biol Buccale* 1975;3:13–28.
67. Van der Velden U. Probing force and the relationship of the probe tip to the periodontal tissues. *J Clin Periodontol* 1979;6:106–114.
68. Mombelli A, Graf H. Depth-force pattern in periodontal probing. *J Clin Periodontol* 1986;13:126–130.
69. Mombelli A, Mühle T, Frigg R. Depth-force patterns of periodontal probing. Attachment gain in relation to probing force. *J Clin Periodontol* 1992;19:295–300.
70. Mombelli A, Mühle T, Brägger U. Probing force dependence of attachment level measurement in the assessment of effects of surgical periodontal treatment with or without NSAID. *Acta Med Dent Helv* 1996;1:34–39.
71. Mombelli A, Mühle T, Brägger U, Lang NP, Bürgin WB. Comparison of periodontal and periimplant probing by depth-force pattern analysis. *Clin Oral Implants Res* 1997;8:448–455.
72. Quirynen M, van Steenberghe D, Jacobs R, Schotte A, Darius P. The reliability of pocket probing around screw-type implants. *Clin Oral Implants Res* 1991;2:186–192.
73. Buser D, Weber HP, Lang NP. Tissue integration of non-submerged implants. 1-year results of a prospective study with 100 ITI hollow-cylinder and hollow-screw implants. *Clin Oral Implants Res* 1990;6:55–61.
74. Adell R, Lekholm U, Rockler B, Brånemark P-I. A 15-year study of osseointegrated implants in the treatment of the edentulous jaw. *Int J Oral Surg* 1981;10:387–416.
75. Cox JF, Zarb GA. The longitudinal clinical efficacy of osseointegrated implants: A 3-year report. *Int J Oral Maxillofac Implants* 1987;2:91–100.
76. Ericsson I, Lekholm U, Brånemark P-I, Lindhe J, Glantz P-O, Nyman S. A clinical evaluation of fixed-bridge restorations supported by the combination of teeth and osseointegrated titanium implants. *J Clin Periodontol* 1986;13:307–312.
77. Lekholm U, Adell R, Lindhe J, et al. Marginal tissue reactions at osseointegrated titanium fixtures. (II) A cross-sectional retrospective study. *Int J Oral Maxillofac Surg* 1986;15:53–61.
78. Nishimura K, Itoh T, Takaki K, Hosokawa R, Naito T, Yokota M. Periodontal parameters of osseointegrated dental implants. A 4-year controlled follow-up study. *Clin Oral Implants Res* 1997;8:272–278.
79. Bauman GR, Mills M, Rapley JW, Hallmon WH. Clinical parameters of evaluation during implant maintenance. *Int J Oral Maxillofac Implants* 1992;7:220–227.
80. Salcetti JM, Moriarty JD, Cooper LF, et al. The clinical, microbial and host response characteristics of the failing implant. *Int J Oral Maxillofac Implants* 1997;12:32–42.
81. Spiekermann H, Jansen VK, Richter EJ. A 10-year follow-up study of IMZ and TPS implants in edentulous mandibles using bar-retained overdentures. *Int J Oral Maxillofac Implants* 1995;10:373–378.
82. Behneke A, Behneke N, d'Hoedt B, Wagner W. Hard and soft tissue reactions to ITI screw implants: 3-year longitudinal results of a prospective study. *Int J Oral Maxillofac Implants* 1997;12:749–757.
83. Brägger U, Bürgin WB, Hämmerle CHF, Lang NP. Associations between clinical parameters assessed around implants and teeth. *Clin Oral Implants Res* 1997;8:412–421.
84. Brägger U, Hugel-Pisoni C, Bürgin WB, Buser D, Lang NP. Correlations between radiographic, clinical and mobility parameters after loading of oral implants with fixed partial dentures. *Clin Oral Implants Res* 1996;7:230–239.
85. Christensen MM, Joss A, Lang NP. Reproducibility of automated periodontal probing around teeth and osseointegrated oral implants. *Clin Oral Implants Res* 1997;8:455–464.
86. Taylor AC, Campbell MM. Reattachment of gingival epithelium to the tooth. *J Periodontol* 1972;43:281–293.
87. Etter TH, Håkanson IH, Lang NP, Trejo PM, Caffesse RG. Healing after standardized clinical probing of the peri-implant soft tissue seal. *Clin Oral Implants Res* 2002;13:571–580.
88. Lang NP, Loe H. The relationship between the width of keratinized gingiva and gingival health. *J Periodontol* 1972;43:623–627.
89. Miyasato M, Crigger M, Egelberg J. Gingival condition in areas of minimal and appreciable width of keratinized gingiva. *J Clin Periodontol* 1977;4:200–209.
90. Strub JR, Gaberthüel TW, Grunder U. The role of attached gingiva in the health of periimplant tissue in dogs. 1. Clinical findings. *Int J Periodontics Restorative Dent* 1991;11:317–333.
91. Warrar K, Buser D, Lang NP, Karring T. Plaque-induced periimplantitis in the presence or absence of keratinized mucosa. An experimental study in monkeys. *Clin Oral Implants Res* 1995;6:131–138.
92. Hanisch O, Cortella CA, Boskovic MM, James RA, Slots J, Wikesjö UM. Experimental peri-implant tissue breakdown around hydroxyapatite-coated implants. *J Periodontol* 1997;68:59–66.
93. Block MS, Gardiner D, Kent JN, Misiak DJ, Finger IM, Guerra L. Hydroxyapatite-coated cylinder implants in the posterior mandible: 10-year observations. *Int J Oral Maxillofac Implants* 1996;11:626–633.
94. Albrektsson T, Zarb GA, Worthington DP, Eriksson R. The long-term efficacy of currently used dental implants. A review and proposed criteria of success. *Int J Oral Maxillofac Implants* 1986;1:11–25.
95. Bengazi F, Wennström JL, Lekholm U. Recession of the soft tissue margin at oral implants. *Clin Oral Implants Res* 1996;7:303–310.
96. Wennström JL, Bengazi F, Lekholm U. The influence of the masticatory mucosa on the peri-implant soft tissue condition. *Clin Oral Implants Res* 1994;5:1–8.
97. Zitzmann NU, Schärer P, Marinello CP. Long-term results of implants treated with guided bone regeneration: A 5-year prospective study. *Int J Oral Maxillofac Implants* 2001;16:355–366.
98. Offenbacher S, Collins JG, Heasman PA. Diagnostic potential of host response mediators. *Adv Dent Res* 1993;7:175–181.
99. Niimi A, Ueda M. Crevicular fluid in the osseointegrated implant sulcus: A pilot study. *Int J Oral Maxillofac Implants* 1995;10:434–436.

100. Boutros SM, Michalowicz BS, Smith QT, Aeppli DM. Crevicular fluid enzymes from endosseous dental implants and natural teeth. *Int J Oral Maxillofac Implants* 1996;11:322–330.
101. Eley BM, Cox SW, Watson RM. Protease activities in periimplant sulcus fluid from patients with permucosal osseointegrated dental implants. Correlation with clinical parameters. *Clin Oral Implants Res* 1991;2:62–70.
102. Apse P, Ellen RP, Overall CM, Zarb GA. Microbiota and crevicular fluid collagenase activity in the osseointegrated dental implant sulcus: A comparison of sites in edentulous and partially edentulous patients. *J Periodontol* 1989;24:96–105.
103. Ataoglu H, Alptekin NO, Haliloglu S, et al. Interleukin 1-beta, tumor necrosis factor-alpha levels and neutrophil elastase activity in peri-implant crevicular fluid. *Clin Oral Implants Res* 2002;13:470–476.
104. Giannopoulou C, Bernard JP, Buser D, Carrel A, Belser UC. Effect of intracrevicular restoration margins on peri-implant health: Clinical, biochemical and microbiological findings around esthetic implants up to 9 years. *Int J Oral Maxillofac Implants* 2003;18:173–181.
105. Hultin M, Gustafsson A, Hallström H, Johansson L-Å, Ekfeldt A, Klinge B. Microbiological findings and host response in patients with periimplantitis. *Clin Oral Implants Res* 2002;13:349–358.
106. Ingman T, Kononen M, Siirila HS, Suomalainen K, Sorsa T. Collagenase, gelatinase and elastase activities in sulcular fluid of osseointegrated implants and natural teeth. *J Clin Periodontol* 1994;21:301–307.
107. Plagnat D, Giannopoulou C, Carrel A, Bernard JP, Mombelli A, Belser UC. Elastase, alpha2-macroglobulin and alkaline phosphatase in crevicular fluid from implants with and without periimplantitis. *Clin Oral Implants Res* 2002;13:227–233.
108. Rühling A, Jepsen S, Kocher T, Plagmann H-C. Longitudinal evaluation of aspartate aminotransferase (AST) in the crevicular fluid of implants with bone loss and signs of progressive disease. *Int J Oral Maxillofac Implants* 1999;14:428–435.
109. Beck CB, Embery G, Langley MS, Waddington RJ. Levels of glycosaminoglycans in periimplant sulcus fluid as a means of monitoring bone response to endosseous dental implants. *Clin Oral Implants Res* 1991;2:179–185.
110. Smedberg JI, Beck CB, Embery G. Glycosaminoglycans in periimplant sulcus fluid from implants supporting fixed or removable prostheses. *Clin Oral Implants Res* 1993;4:137–143.
111. Kao RT, Curtis DA, Richards DW, Preble J. Increased interleukin-1beta in the crevicular fluid of diseased implants. *Int J Oral Maxillofac Implants* 1995;10:696–701.
112. Esposito M, Thomsen P, Mölne J, Gretzer C, Ericson LE, Lekholm U. Immunohistochemistry of soft tissues surrounding late failures of Brånemark implants. *Clin Oral Implants Res* 1997;8:352–366.
113. Gualini F, Berglundh T. Immunohistochemical characteristics of inflammatory lesions at implants. *J Clin Periodontol* 2003;30:14–18.
114. Piattelli A, Scarano A, Piattelli M. Histologic observations on 230 retrieved dental implants: 8 years' experience (1989–1996). *J Periodontol* 1998;69:178–184.
115. Sanz M, Alandez J, Lazaro P, Calvo JL, Quirynen M, van Steenberghe D. Histo-pathologic characteristics of periimplant soft tissues in Brånemark implants with 2 distinct clinical and radiological patterns. A histometric and ultrastructural study. *Clin Oral Implants Res* 1991;2:128–134.
116. Brånemark P-I, Hansson BO, Adell R, et al. Osseointegrated implants in the treatment of the edentulous jaw. Experience from a 10-year period. *Scand J Plast Reconstr Surg* 1977;16(suppl):1–132.
117. Lekholm U, van Steenberghe D, Herrmann I, et al. Osseointegrated implants in the treatment of partially edentulous jaws: A prospective 5-year multicenter study. *Int J Oral Maxillofac Implants* 1994;9:627–635.
118. Gröndahl K, Lekholm U. The predictive value of radiographic diagnosis of implant instability. *Int J Oral Maxillofac Implants* 1997;12:59–64.
119. Worthington P, Bolender CL, Taylor TD. The Swedish system of osseointegrated implants: Problems and complications encountered during a 4-year trial period. *Int J Oral Maxillofac Implants* 1987;2:77–84.
120. Olivé J, Aparicio C. The Periotest method as a measure of osseointegrated oral implant stability. *Int J Oral Maxillofac Implants* 1990;5:390–400.
121. Naert I, Rosenberg D, van Steenberghe D, Tricio JA, Nys M. The influence of splinting procedures on the periodontal and peri-implant tissue damping characteristics. A longitudinal study with the Periotest device. *J Clin Periodontol* 1995;22:703–708.
122. Teerlinck J, Quirynen M, Darius P, van Steenberghe D. Periotest: An objective clinical diagnosis of bone apposition toward implants. *Int J Oral Maxillofac Implants* 1991;6:55–61.
123. Aparicio C. The use of the Periotest value as the initial success criteria of an implant: 8-year report. *Int J Periodontics Restorative Dent* 1997;6:151–161.
124. van Steenberghe D, Tricio J, Naert I, Nys M. Damping characteristics of bone-to-implant interfaces. A clinical study with the Periotest device. *Clin Oral Implants Res* 1995;6:31–39.
125. Meredith N. Assessment of implant stability as a prognostic determinant. *Int J Prosthodont* 1998;11:491–501.
126. Meredith N, Alleyne D, Cawley P. Quantitative determination of the stability of the implant-tissue interface using resonance frequency analysis. *Clin Oral Implants Res* 1996;7:261–266.
127. Friberg B, Sennerby L, Meredith N, Lekholm U. A comparison between cutting torque and resonance frequency measurements of maxillary implants. A 20-month clinical study. *Int J Oral Maxillofac Surg* 1999;28:297–303.
128. Heo SJ, Sennerby L, Odersjö M, Grånström G, Tjellström A, Meredith N. Stability measurements of craniofacial implants by means of resonance frequency analysis. A clinical pilot study. *J Laryngol Otol* 1998;112:537–542.
129. Meredith N, Book K, Friberg B, Jemt T, Sennerby L. Resonance frequency measurements of implant stability in vivo. A cross-sectional and longitudinal study of resonance frequency measurements on implants in the edentulous and partially dentate maxilla. *Clin Oral Implants Res* 1997;8:226–233.
130. Meredith N, Shagaldi F, Alleyne D, Sennerby L, Cawley P. The application of resonance frequency measurements to study the stability of titanium implants during healing in the rabbit tibia. *Clin Oral Implants Res* 1997;8:234–243.
131. Rasmusson L, Meredith N, Sennerby L. Measurements of stability changes of titanium implants with exposed threads subjected to barrier membrane induced bone augmentation. An experimental study in the rabbit tibia. *Clin Oral Implants Res* 1997;8:316–322.

132. Rasmusson L, Kahnberg KE, Tan A. Effects of implant design and surface on bone regeneration and implant stability: An experimental study in the dog mandible. *Clin Implant Dent Relat Res* 2001;3:2–8.
133. Friberg B, Sennerby L, Linden B, Gröndahl K, Lekholm U. Stability measurements of one-stage Brånemark implants during healing in mandibles. A clinical resonance frequency analysis study. *Int J Oral Maxillofac Surg* 1999;28:266–272.
134. Barewal RM, Oates T, Meredith N, Cochran DL. Resonance frequency measurement of implant stability in vivo on implants with a sandblasted and acid-etched surface. *Int J Oral Maxillofac Implants* 2003;18:641–651.
135. Smith DE, Zarb GA. Criteria for success of osseointegrated endosseous implants. *J Prosthet Dent* 1989;62:567–572.
136. Feloutzis A, Lang NP, Tonetti MS, et al. IL-1 gene polymorphism and smoking as risk factors for periimplant bone loss in a well-maintained population. *Clin Oral Implants Res* 2003;14:10–17.
137. Brägger U, Pasquali L, Rylander H, Carnes D, Kornman KS. Computer-assisted densitometric image analysis in periodontal radiography. *J Clin Periodontol* 1988;15:27–37.
138. Lang NP, Hill RW. Radiographs in periodontics. *J Clin Periodontol* 1977;4:16–28.
139. Sewerin IP, Gottfredsen K, Stoltze K. Accuracy of radiographic diagnosis of periimplant radiolucencies—An in vitro experiment. *Clin Oral Implants Res* 1997;8:299–304.
140. Wennström JL, Palmer RM. Consensus report of session C. In: Lang NP, Karring T, Lindhe J (eds.). *Proceedings of the 3rd European Workshop on Periodontology*. Berlin: Quintessenz, 1999:255–259.
141. Brägger U, Pasquali L, Weber HP, Kornman KS. Computer-assisted densitometric image analysis (CADIA) for the assessment of alveolar bone density changes. *J Clin Periodontol* 1989;16:46–52.
142. Fourmoussis I, Brägger U. Radiologic interpretation of peri-implant structures. In: Lang NP, Karring T, Lindhe J (eds.). *Proceedings of the 3rd European Workshop on Periodontology*. Berlin: Quintessenz, 1999:228–241.